

AN ASTROMETRIC COMPANION TO THE NEARBY METAL-POOR, LOW-MASS STAR LHS 1589.^{1,2}

SÉBASTIEN LÉPINE³, R. MICHAEL RICH⁴, MICHAEL M. SHARA³, KELLE L. CRUZ^{3,5}, AND ANDREW SKEMER⁶

Submitted to the Astrophysical Journal

ABSTRACT

We report the discovery of a companion to the high proper motion star LHS 1589, a nearby high-velocity, low-mass subdwarf. The companion (LHS 1589B) is located $0.224 \pm 0.004''$ to the southwest of the primary (LHS 1589A), and is 0.5 magnitude fainter than the primary in the K_s band. The pair was resolved with the IRCAL infrared camera at Lick Observatory, operating with the Laser Guide Star Adaptive Optics system. A low-resolution spectrum of the unresolved pair obtained at the MDM observatory shows the source to be consistent with a cool subdwarf of spectral subtype sdK7.5. A photometric distance estimate places the metal-poor system at a distance $d = 81 \pm 18$ parsecs from the Sun. We also measure a radial velocity $V_{rad} = 67 \pm 8 \text{ km s}^{-1}$ which, together with the proper motion and estimated distance, suggests that the pair is roaming the inner Galactic halo on a highly eccentric orbit. With a projected orbital separation $s = 18.1 \pm 4.8 \text{ AU}$, and a crude estimate of the system's total mass, we estimate the orbital period of the system to be in the range $75 \text{ yr} < P < 500 \text{ yr}$. This suggests that the dynamical mass of the system could be derived astrometrically, after monitoring the orbital motion over a decade or so. The LHS 1589AB system could thus provide a much needed constraint to the mass-luminosity relationship of metal-poor, low-mass stars.

Subject headings: binaries: close — techniques: high angular resolution — stars: low-mass, brown dwarfs — stars: subdwarfs — stars: Population II — Galaxy: solar neighborhood

1. INTRODUCTION

Astrometric binaries, defined as physical pairs whose projected orbital motion can be mapped out, are objects of high interest in astrophysics because the gravitational masses of the components can be directly calculated. Such systems are critical in constraining the mass-luminosity relationship (MLR), one of the most fundamental relationships in astrophysics (Malkov *et al.* 1997; Henry *et al.* 1999; Ségransan, D. *et al.* 2000; Delfosse *et al.* 2000; Martinache *et al.* 2007). The MLR allows one to convert the luminosity function (LF), an observable quantity, into the mass function or initial mass function (IMF) which is fundamental in understanding stellar populations (D'Antona 1998). The MLR is particularly critical to obtain reliable estimates of the stellar mass function and baryonic content of the Galaxy (Chabrier & Méra 1997; Phleps *et al.* 2000). The MLR further provides constraints to stellar evolutionary models (Baraffe *et al.* 1998).

Differences have been observed in the luminosity functions of open and globular clusters which are best explained by a dependence of the MLR on metallicity (von Hippel *et al.* 1996). In low-mass stars, particularly the M dwarfs/subdwarfs, the spectral energy distribution, dominated by molecular bands of metal oxides and hydrides, is strongly affected by variations in metallicity (Allard & Hauschildt 1995). A high metal content yields stronger molecular bands,

which shift the energy output from the optical to the infrared, making the star redder. Assuming the bolometric luminosity to be simply dependent on stellar mass, the MLR for metal-poor stars is thus expected to be shifted to brighter magnitudes in the optical, and possibly also to fainter magnitudes in the infrared. Theoretical models combined with observational evidence suggest that metallicity effects on the MLR are more pronounced in the visual bands than in the IR (Delfosse *et al.* 2000), at least for M dwarfs of near-solar metallicity ($-0.5 < \log[Fe/H] < 0.0$). It has been suggested that the MLR can be assumed to be independent of metallicity in the K band (Bonfils *et al.* 2005), although this remains to be demonstrated for low-mass stars over a broad range of Fe/H values. It is thus critical to verify the MLR calibration for stars over a wider range of metallicities, particularly for very metal-poor ($\log[Fe/H] < -1.0$) M subdwarfs.

There is extensive literature on the topic of low-mass astrometric doubles and the calibration of the MLR for low-mass dwarfs of near-solar metallicity (Population I). Early calibrations, in the classic papers by Liebert & Probst (1987) and Henry & McCarthy (1993), relied on relatively few systems with well determined orbits and masses. Other known double stars have since had their orbits monitored with speckle interferometry (Mason *et al.* 1999; Woitas *et al.* 2000), or with the Fine Guidance Sensor on HST (Franz *et al.* 1998; Henry *et al.* 1999; Torres *et al.* 1999; Benedict *et al.* 2000, 2001), yielding accurate masses for both components. New close doubles have also been resolved among nearby disk dwarfs with the NICMOS camera on HST (Pravdo *et al.* 2004), with adaptive optics (AO) observations from CFHT (Delfosse *et al.* 1999b), with AO masking interferometry from Palomar (Lloyd *et al.* 2006), and with Laser Guide Star Adaptive Optics (LGS AO) observations from Keck (Pravdo *et al.* 2006).

In contrast, there exists no calibration of the mass-luminosity relationship for the more metal-poor, low-mass subdwarfs (Population II). Perhaps the chief reason for this is that most metal-poor stars in the vicinity of the Sun are kinematically associated with the Galactic halo population, and

¹ Based on observations performed with the Laser Guide Star Adaptive Optics system at the Lick Observatory, operated by the University of California system.

² Based on observations conducted at the MDM observatory, operated jointly by the University of Michigan, Dartmouth College, the Ohio State University, Columbia University, and the University of Ohio.

³ Department of Astrophysics, Division of Physical Sciences, American Museum of Natural History, Central Park West at 79th Street, New York, NY 10024, USA

⁴ Department of Astrophysics, University of California at Los Angeles, Los Angeles, CA 90095, USA, rmr@astro.ucla.edu

⁵ NSF Astronomy and Astrophysics Postdoctoral Fellow

⁶ Department of Astronomy and Steward Observatory, University of Arizona, 933 N Cherry Ave., Tucson, AZ 85721, USA, askemer@as.arizona.edu




FIG. 1.— Multi-epoch chart of the high proper motion star LHS 1589 generated from the Digitized Sky Surveys. The field is $4.25'$ on the side. Left: first Palomar Sky Survey image (POSS-I) circa 1954.0. Right: Second Palomar Sky Survey image (POSS-II) circa 1988.9. The motion of LHS 1589 is quite apparent. The box shows the location of the field imaged with the Lick Adaptive Optics system on October 2006.

are thus relatively scarce. Most studies estimate there is one halo star in the Solar Neighborhood for each 200-300 disk stars, which means that one expects to find only ~ 20 halo stars within 25 parsecs of the Sun. The MLR for low-mass, Pop II subdwarfs has also been lacking because of the difficulty in measuring accurate metallicities in low-mass stars, although significant progress has recently been made on that front (Bean *et al.* 2006; Woolf & Wallerstein 2006).

In any case, the existence of wide binaries among the halo population has been demonstrated. Positive identification of low-mass subdwarfs in lists of common proper motion doubles (Chanamé & Gould 2004) suggests that the binary frequency among old metal-poor stars may be as high as that of the local disk population. Recently, a companion was resolved only $0.15''$ from the nearby ($\approx 100pc$) metal-poor sdG2 subdwarf Ross 530 (Law *et al.* 2006). If they could be found, subdwarf doubles with components of lower masses (sdK/sdM), and with separations short enough for the orbital motion to be astrometrically monitored, would be of great value to astronomy.

We have recently undertaken a massive study of the local population of metal-poor, low-mass stars, following the identification of thousands of high-velocity subdwarfs in the new LSPM-north catalog of stars with proper motions $\mu > 0.15'' \text{ yr}^{-1}$ (Lépine & Shara 2005) and in its upcoming southern-sky complement, the LSPM-south (Lépine *et al.* 2007, in preparation). Here, we report the discovery of a close companion to the high proper motion star LHS 1589, which our spectroscopic observations confirm to be a metal-poor, cool subdwarf. In §2, we describe our Adaptive Optics observations, performed at Lick Observatory with the Laser Guide Star system. Our spectroscopic observations are detailed in §3. We discuss in §4 the prospects for using this system to constrain the mass-luminosity relationship in the metal-poor, low-mass star regime.

2. ADAPTIVE OPTICS OBSERVATIONS

From the LSPM-north catalog (Lépine & Shara 2005), we have assembled a list of probable low-mass subdwarfs with estimated (photometric) distances $d < 100pc$ from the Sun. A first subset of 18 targets were observed on the nights of




FIG. 2.— Adaptive optics image of LHS 1589, resolving this high proper motion star into two distinct sources. The secondary is found $0.224'' \pm 0.004''$ to the southwest of the primary. The image shown here represents only the central part of the full IRCAL field, as was constructed from a set of five dithered frames.

2006 September 16-17, with the Shane 3.0-meter telescope at Lick Observatory on Mount Hamilton. Adaptive Optics (AO) observations were performed under operation of the Laser Guide Star (LGS) system. High angular resolution images were recorded with the IRCAL infrared camera, a 256×256 pixels Rockwell PICNIC HgCdTe array (Lloyd *et al.* 2000). The camera has an effective field of view of approximately $19.4'' \times 19.4''$ in the focal plane. It is however slightly anamorphic, with a measured plate scale of $0.0754'' \text{ pixel}^{-1}$ in the direction of Right Ascension, and $0.0780'' \text{ pixel}^{-1}$ in the direction of Declination (Perrin 2006). We operated the laser guide star system using the science target as a reference tip/tilt star. A total of 5 images were obtained in the K band with a 5 point dithering pattern.

No companions were detected in 17 of the 18 targets. These stars yielded radially symmetric images, with the first Airy ring clearly visible. We conclude that these stars are either single star systems, or that they are multiples unresolved to within the diffraction limit of the telescope.

One of our targets, the high proper motion star LHS 1589

(see Figure 1), was however resolved into a pair of close point sources. All five images from the dithering pattern showed the same pair of resolved point sources, with similar separation and orientation. A composite image was built using the *drizzle* algorithm in the IRAF STSDAS package, halving the effective pixel scale to $0.0377'' \times 0.0390''$. Figure 2 shows the central $3.5'' \times 3.5''$ part of the composite image. Point spread function fitting of the two resolved sources yields an angular separation $\rho = 0.224'' \pm 0.004''$, with a position angle on the sky from component A to B of $\phi = 206.3 \pm 1.3$ degrees.

We can rule out with near certainty the possibility that the two stars are a chance alignment. Each component is bright enough to be detected in any of the POSS-I or POSS-II images, but scans from the Digitized Sky Surveys (DSS) covering the piece of sky imaged by IRCAL is clearly devoid of any optical source down to the magnitude limit of the photographic plates (see Fig.1). Likewise, an image from the 2MASS data archive shows a single, unresolved source near the location of LHS 1589, though the 2MASS data was acquired in 1999, at a time when LHS 1589 was $> 5''$ away from its current location. Any background source located near the current location of LHS 1589 would have been clearly resolved in the 2MASS image. Hence, the two point sources are most likely co-moving stars. The only alternative would be for component B to be a strongly variable or transient source that just happened to erupt within a short angular distance of LHS 1589 at the time of the AO imaging, a very unlikely event.

Component B currently lies to the South-West of the brighter component A. Based on our aperture photometry, the secondary is 0.52 magnitudes fainter than the primary in the K_s band, which we can write down as $\Delta K_s = (K_s)_B - (K_s)_A = 0.52$. Infra-red 2MASS photometry of the unresolved source yields a total magnitude $(K_s)_{A+B} = 11.33$ mag. We can derive magnitudes for the individual components using:

$$(K_s)_A = (K_s)_{A+B} + 2.5 \log 1 + 10^{-\Delta K_s/2.5}, \quad (1)$$

from which we calculate $K_s = 11.85$ mag for component A, and $K_s = 12.36$ mag for component B.

3. SPECTROSCOPIC OBSERVATIONS

A red optical spectrum of the unresolved binary system was collected at the MDM observatory with the 1.3m McGraw-Hill telescope on the night of 14 October 2006. We used the Mark III spectrograph with the 300 lines/mm grating blazed at $8000\text{\AA}$. The spectrum was recorded with a LORAL 2048×2048 CCD camera ("Wilbur"), yielding a spectral resolution of $3.11\text{\AA}$ per pixel. Reduction was done with IRAF, including standard flat-fielding, sky-subtraction, extraction, calibration, and telluric correction. Wavelength calibration was determined from a NeArXe arc spectrum collected right after the exposure on the star, yielding an estimated calibration accuracy of $\pm 0.35\text{\AA}$. Flux calibration was determined based on observations of the spectrophotometric standard star Hiltner 600.

The reduced, composite spectrum is displayed in Figure 3. The relatively weak TiO/CaH band ratio is consistent with a cool subdwarf of spectral type sdK-sdM. We calculated the three spectroscopic indices CaH2, CaH3, and TiO5, which measure the depth of the CaH and TiO molecular bandheads near $7040\text{\AA}$ (Reid, Hawley, & Gizis 1995), finding $\text{CaH2} = 0.781$, $\text{CaH3} = 0.890$, $\text{TiO5} = 0.906$. In the revised classification system of Lépine, Rich, & Shara (2007), these values yield a




FIG. 3.— Composite spectrum of the halo subdwarf binary LHS 1589AB. The strong CaH molecular band near $7000\text{\AA}$ flanked by a relatively weak TiO bands indicates that the star is a metal-poor, low-mass subdwarf. Based on the strength of the CaH band, the spectral subtype of the composite spectrum is sdK7.5. Two other stars from our MDM spectroscopic survey are shown for comparison: the K7.5 dwarf LSPM J2328+3219, and the esdK7.5 extreme subdwarf LSPM J1320+1129.

spectral type sdK7.5. The TiO/CaH bandstrength ratio, which is the main metallicity diagnostic in cool subdwarfs, is quite weak in LHS 1589AB, and the star actually falls just short of being classified an extreme subdwarf (esdK7.5).

A direct comparison with spectra from other cool dwarfs and subdwarfs confirms this impression. Figure 3 shows MDM spectra of the high proper motion stars LSPM J2328+3219 and LSPM J1320+1129, respectively classified as K7.5 and esdK7.5, respectively. Each is typical of stars of its own metallicity subclass, and both have CaH bandstrengths similar to those observed in LHS 1589AB. The TiO bands in LHS 1589 are clearly intermediate between the two comparison objects, but are closer to those of the esdK7.5 star, which again indicates that LHS 1589AB consists of two subdwarfs (sd) close to the extreme subdwarf (esd) metallicity class.

A radial velocity was calculated from the shift of individual lines in the K II doublet and Ca II triplet, whose centroids were calculated in IRAF with the SPECRED package. With a 1-pixel resolution of $3.11\text{\AA}$ and signal-to-noise ratio of 30, we estimate that we can measure radial velocities to $\approx 8\text{ km/s}$. After correction for the Earth's orbital motion, we determine a heliocentric radial velocity $V_{\text{rad}} = +67 \pm 8\text{ km/s}$.

4. DISCUSSION

The star LHS 1589 was first discovered as a high proper motion star in the Lowell Proper motion survey (Giglas, Burnham, & Thomas 1961), and was initially known




FIG. 4.— Color-magnitude diagram of nearby stars with measured trigonometric parallaxes, from various sources. Spectroscopically confirmed subdwarfs (sdM) are shown as filled triangles; spectroscopically confirmed extreme subdwarfs (esdM) are shown as open circles. Component A in LHS 1589 has an estimated color $V-K_s=3.5$ (vertical line). Based on our spectroscopy (Fig.3), we assume that LHS 1589 is a subdwarf located close to the extreme subdwarfs, from which we estimate an absolute magnitude $M_{K_s} \approx 7.3 \pm 0.5$ (horizontal lines). This in turn suggests a distance $d = 81 \pm 18$ pc for the system.

as G 79-69. The star was later included in the Luyten Half Second catalog (Luyten 1979) under the catalog number 1589. Olin J. Eggen also included the object in his catalog of stars with proper motions $\mu > 0.7'' \text{ yr}^{-1}$ under the designation PM 03436+1134. Eggen also recognized it to be a metal-poor subdwarf based on its relatively blue $R-I$ color (Eggen (1983).

It is critical that the distance to the pair be properly measured so that we can estimate its projected physical separation and probable orbital period. This in turn will determine the potential for the system to have its orbital motion mapped out astrometrically. We have compiled published trigonometric parallaxes of spectroscopically confirmed, nearby subdwarfs (Monet *et al.* 1992; van Altena *et al.* 1995). Figure 4 shows the resulting $[M_{K_s}, V-K_s]$ color-magnitude diagram. Optical V magnitudes are from the original parallax tables, while infra-red K_s magnitudes are from the 2MASS All-Sky Point Source catalog (Cutri *et al.* 2003). Spectroscopically confirmed subdwarfs (sdK, sdM) and extreme subdwarfs (esdK, esdM) are plotted with distinct symbols in Fig. 4 (filled triangles and open circles, respectively). The list includes subdwarfs classified by Gizis (1997) and Reid & Gizis (2005), as well as subdwarfs classified in our own spectroscopic survey (Lépine *et al.* 2007, in preparation).

As can be seen in Fig.4, the color-magnitude relationship has a strong dependence on metallicity for stars with $3 < V-K_s < 5$, with more metal-poor stars being less luminous at a given color. The parallax data suggest that $\Delta M_{K_s} \sim 1.7 \Delta(V-K_s)$ for sdK/sdM subdwarfs. Our own measurements give $(K_s)_B - (K_s)_A = 0.52$, from which we estimate that $\Delta(V-K_s) = (V-K_s)_B - (V-K_s)_A = 0.31$, which yields $\Delta V = V_B - V_A = 0.83$. This suggests that component B is about 2.15 times fainter in the V band than component A. The inte-

TABLE 1
SUMMARY OF DATA ON LHS 1589AB

Observable	value
Primary R.A. (J2000) ^a	03 46 24.53
Primary Decl. (J2000) ^a	+11 43 12.0
μ R.A. ($'' \text{ yr}^{-1}$)	0.783
μ Decl. ($'' \text{ yr}^{-1}$)	-0.178
V_{A+B} (mag)	14.93
J_{A+B} (mag)	12.06
H_{A+B} (mag)	11.51
$(K_s)_{A+B}$ (mag)	11.33
$(K_s)_B - (K_s)_A$ (mag)	0.52
composite spectral type	sdK7.5
V_{rad} (km s^{-1})	$+67 \pm 8$
Angular separation ($''$)	0.224 ± 0.004
position angle (degrees)	206.3 ± 1.3
Derived quantity	value
V_A (mag)	15.35
$(K_s)_A$ (mag)	11.85
V_B (mag)	16.18
$(K_s)_B$ (mag)	12.36
photometric distance (pc)	81 ± 18
U (km s^{-1})	-155 ± 8
V (km s^{-1})	-240 ± 16
W (km s^{-1})	$+116 \pm 9$
m_A ($M_\odot$)	≈ 0.25
m_B ($M_\odot$)	≈ 0.30

^a Positions are given at the 2000.0 epoch and for the J2000 equinox.

grated V band magnitude of the system is $V_{A+B}=14.93$, as measured by Harrington & Dahn (1980). We can use Equation 1 again to calculate the V magnitudes of components A and B, substituting V for K_s in the equation. We find $V_A = 15.35$ for the primary, and $V_B = 16.18$ for the secondary. This in turn suggests optical-to-infrared colors $(V-K_s)_A = 3.5$ and $(V-K_s)_B = 3.8$.

Figure 4 displays our best guess at the absolute magnitude of LHS 1589A. If we assume $V-K_s \approx 3.5$, and further assume the star to be near the limit between sdK-sdM and esdK-esdM subdwarfs, then we find that LHS 1589A has an absolute magnitude $M_{K_s} = 7.3 \pm 0.5$. An independent estimate of the absolute magnitude can be obtained based on the star's spectrum. Relationships between the spectral subtype and absolute magnitude have been calibrated by Lépine, Rich, & Shara (2003) and Reid & Gizis (2005) for groups of dwarfs, subdwarfs, and extreme subdwarfs. For an sdK7.5 subdwarf, just halfway between the subdwarf and extreme subdwarf classes, the subtype-luminosity relationships are consistent with an absolute magnitude $M_{K_s} \approx 6.9$, which agrees quite well with the color-magnitude estimate given above.

Based on an absolute magnitude $M_{K_s} = 7.3 \pm 0.5$, we derive a photometric distance $d = 81 \pm 18$ pc. The only recorded parallax for LHS 1589, a relatively crude 2.1 ± 7.9 milliarcsecond (Harrington & Dahn 1980), is consistent with our photometric distance estimate at the $\approx 1.3\sigma$ level. At a distance of 81 pc, and assuming $V_{rad} = +67 \text{ km s}^{-1}$, the system would have components of motion $(U, V, W) = (-155, -240, +117) \text{ km s}^{-1}$ in the local standard of rest. An integration in the Galactic potential yields a very eccentric orbit sharply inclined to the Galactic plane, consistent with a membership with the inner Galactic Halo.

Is it also possible to estimate the mass of the system, with the caveat that existing mass-luminosity and mass-color rela-

tionships have all been calibrated using dwarf stars of roughly solar metallicity, while the components of LHS 1589 are really metal-poor subdwarfs. Both the mass-luminosity relationship in the V band and the mass-color ($V - K_s$) relationship are significantly dependent on metallicity (Bonfils *et al.* 2005; Delfosse *et al.* 2000), and thus cannot be used. The mass-luminosity relationship in the infrared K band, however, does appear to be nearly independent on metallicity. According to the Delfosse *et al.* (2000) calibration, LHS1589A, with $M_K = 7.3 \pm 0.5$ should have a mass in the range $0.2M_\odot < m_A < 0.3M_\odot$. Component B, with $M_K = 7.8 \pm 0.5$, should have a mass in the range $0.15M_\odot < m_B < 0.25M_\odot$. The total mass of the system should thus be roughly in the range of $0.35M_\odot < m_{A+B} < 0.55M_\odot$.

At first glance, these mass estimates seem suspiciously low. This is because mass to spectral-subtype relationships for dwarf stars indicate that a star of spectral subtype K7.5 should have a mass of $\approx 0.6 M_\odot$ (Baraffe & Chabrier 1996; Henry 2004). If one assumes that the mass to spectral-subtype relationship applies similarly to the subdwarfs, then this would suggest a mass $\approx 0.6 M_\odot$ for LHS 1589A, clearly at odds with our the $0.2\text{--}0.3M_\odot$ estimate. However, one should note that this mass to spectral-subtype relationship has only been calibrated for K and M dwarfs of roughly solar metallicity, and has not been verified for sdK/sdM subdwarfs. In fact, spectral subtypes are usually a measure of the effective temperature, and subdwarfs are expected to be both smaller and hotter than dwarfs of comparable masses. This suggests that a subdwarf of a given subtype is indeed likely to be less massive than a dwarf of similar subtype. In fact, if we assume that M_{K_s} is dependent only on the mass of the star and not its metallicity, as suggested by Delfosse *et al.* (2000), then the subtype-luminosity relationships of (Lépine, Rich, & Shara 2003) and (Reid & Gizis 2005) indicate that an sdK7.5 star like LHS 1589A should have roughly the same mass as an M3-M4 dwarf, which itself has a mass of $\approx 0.3 M_\odot$, a value consistent with our mass estimate.

It is also possible that the mass-luminosity relationship in the K band might not be as tight as suggested in Delfosse *et al.* (2000). It would make sense for subdwarfs to be slightly underluminous in the K band, to compensate for the fact that they are overluminous in the optical, and assuming that stars

of a given mass all have the same bolometric luminosities. In which case, the current mass- M_{K_s} relationships, calibrated on K-M dwarfs, would indeed underestimate the mass of a subdwarf of the same M_{K_s} . The only way to resolve this conundrum is to obtain dynamical masses for components of subdwarf binaries.

Assuming for now that the total mass of the system is in the range $0.35 M_\odot < m_{A+B} < 0.55M_\odot$, then with a projected orbital separation $s = 18.1 \pm 4.8$ AU we estimate an orbital period in the range $75 \text{ yr} \lesssim P \lesssim 500 \text{ yr}$. At the lower orbital period regime, a fraction of the orbit could be mapped within a decade, from which the total dynamical mass of the system could be constrained with much greater accuracy (Schaefer *et al.* 2006).

5. CONCLUSIONS

The low-mass subdwarf double LHS 1589AB should be considered a high priority target for high angular resolution, astrometric monitoring. The system is a good candidate for a direct measurement of dynamical masses, and may prove critical for calibrating the mass-magnitude relationship in metal-poor stars. At the very least, an accurate parallax for the system should be obtained, in order to estimate how much of an investment in time would be required to map out the orbital motion. Systems with much shorter orbital separations would clearly be more useful, however. Our demonstration that astrometric doubles do exist *and* can be detected among nearby subdwarfs, provides renewed impetus in the search for additional systems. We are currently expanding our LGS-AO survey with the aim of surveying all known low-mass subdwarfs within 100 parsecs of the Sun.

Acknowledgments

The authors wish to thank the staff at Lick Observatory, in particular Elinor Gates, for their operation of the Laser Guide Star system and for outstanding support. This research program was supported by NSF grant AST-0607757 at the American Museum of Natural History. SL and MMS also gratefully acknowledge support from Mr. Hilary Lipsitz. KLC is supported by an NSF Astronomy and Astrophysics Postdoctoral Fellowship under award AST-040148.

REFERENCES

- Allard, F., & Hauschildt, P. H. 1995, *ApJ*, 445, 433
 Baraffe, I., & Chabrier, G. 1996, *ApJ*, 461, L51
 Baraffe, I., *et al.* 1998, *A&A*, 337, 403
 Bean, J. L. *et al.* 2006, *ApJ*, 652, 1604
 Benedict, F., *et al.* 2000, *AJ*, 120, 1106
 Benedict, F., *et al.* 2001, *AJ*, 121, 1607
 Bonfils, X., *et al.* 2005, *A&A*, 442, 635
 Chabrier, G., & Méra, D. 1997, *A&A*, 328, 83
 Chanamé, J., & Gould, A. 2004, 601, 289
 Cutri, R. M. *et al.* 2003, The 2MASS All-Sky Catalog of Point Sources, University of Massachusetts and Infrared Processing and Analysis Center, (IPAC/California Institute of Technology)
 D'Antona, F. 1998, The Stellar Initial Mass Function, APS Conference Series, Vol. 142, p. 157
 Delfosse, X. *et al.* 1999, *A&A*, 344, 897
 Delfosse, X. *et al.* 2000, *A&A*, 364, 217
 Eggen, O. J. 1983, *ApJS*, 51, 183
 Franz, O. G., *et al.*, 1998, *AJ*, 116, 1432
 Giclas H. L., Burnham, R., & Thomas N.G. 1961, *Lowell Obs. Bull.*, 5, 61
 Gizis, J. E. 1997, *AJ*, 113, 806 (G97)
 Golimowski, D. A., *et al.* 2000, *AJ*, 120, 2082
 Harrington, R. S., & Dahn, C. C. 1980, *AJ*, 85, 454
 Henry, T. J., & McCarthy, D. W. 1993, *AJ*, 106, 773
 Henry, T. J., *et al.* 1999, *ApJ*, 512, 864
 Henry, T. J. 2004, *ASP Conf. Series*, 318, 159
 von Hippel, T., *et al.* 1996, *AJ*, 112, 192
 Law, N. M., Hodgkins, S. T., & Mackay, C. D. 2006, *MNRAS*, 368, 1917
 Lépine, S., Rich, R. M., & Shara, M. M. 2003, *AJ*, *submitted*
 Lépine, S., & Shara, M. M. 2005, *AJ*, 129, 1483
 Lépine, S., Rich, R. M., & Shara, M. M. 2007, *ApJ*, *submitted*
 Liebert, J., & Probst, R. G. 1987, *ARA&A*, 25, 473
 Lloyd, J. P. *et al.* 2000, *Proc. SPIE*, 4008, 814
 Luyten W. J. 1979, LHS Catalogue: a catalogue of stars with proper motions exceeding 0.5" annually, University of Minnesota, Minneapolis (CDS-Vizier catalog number I/87B)
 Malkov, O. Yu., Piskunov, A. E., & Shpil'Kina, D. A. 1997, *A&A*, 320, 79
 Martinache, F. *et al.* 2007, *ApJ*, 661, 496
 Mason, B. D., Douglas, G., & Hartkopf, W. I. 1999, *AJ*, 117, 1023
 Monet, D. G., *et al.* 1992, *AJ*, 103, 638
 Perrin, M. D. 2006, Ph.D. Thesis, UC Berkeley
 Phleps, S., *et al.* 2000, *A&A*, 356, 108
 Pravdo, S. H., *et al.* 2004, *AJ*, 617, 1323
 Lloyd, J. P. *et al.* 2006, *ApJ*, 650, 131
 Pravdo, S., *et al.* 2006, *ApJ*, 649, 389
 Reid, I. N., Hawley, S., & Gizis, J. E. 1995, *AJ*, 110, 1838
 Reid, I. N., & Gizis, J. E. 2005, *PASP*, 117, 676
 Schaefer, G. H., *et al.* 2006, *AJ*, 132, 2618
 Ségransan, D. *et al.* 2000, *A&A*, 364, 665

Torres, G., *et al.* 1999, AJ, 117, 562

Van Altena W.F., Lee J.T., & Hoffleit E.D. 1995, The General Catalogue of Trigonometric Stellar Parallaxes, Fourth Edition, Yale University Observatory

Witas, J., *et al.* 2000, A&A, 353, 253

Wolf, V. M., & Wallerstein, G. 2006, PASP, 118, 218